

Differentiating between Dengue Fever from Other Febrile Illnesses Using Haematological Parameters

P PRIYANKA, US DINESH

ABSTRACT

Introduction: Dengue fever is the most common arthropod borne disease and major public health concerns in India. The clinical presentation of this disease is difficult to distinguish from other febrile illnesses like malaria and typhus fever. Correlation of haematological parameters helps to differentiate these diseases in early stage.

Aim: The present study aimed to identify the haematological features useful for discriminating dengue from other febrile illnesses and to evaluate the accuracy of these haematological parameters.

Materials and Methods: A retrospective study was done to differentiate between dengue and other febrile illnesses between January 2017 to July 2017 at Sri Dharmasthala Manjunatheshwara College of Medical Sciences and Hospital, Dharwad Karnataka. Data regarding

haematological parameters were collected in 170 cases classified as dengue (D) and 170 cases classified as non dengue (ND) based on laboratory tests.

Results: The following parameters were significantly lower in patients with DF as compared to non dengue patients (p -value less than 0.05); WBC, Platelets, Neutrophils, Eosinophils.

The following parameters were significantly higher in patients with DF as compared to non dengue cases ($p < 0.05$): Haemoglobin, PCV, RBC count, Lymphocytes, Monocytes. Multiple logistic regression model used in this study showed two haematological predictors with positive association to confirm dengue: WBC count $< 4000/\text{cumm}$ and platelet count < 1 lakh/cumm (likely to be present in dengue cases).

Conclusion: The study helps to differentiate dengue from other febrile illnesses at an early stage and avoids a large number of other investigation to diagnose dengue from other febrile illness.

Keywords: Febrile illness, Haemoglobin, Lymphocytes, Platelet count

INTRODUCTION

Dengue is a mosquito borne tropical disease presenting with acute febrile illness caused by an arbovirus transmitted by *Aedes aegypti* mosquito [1-5]. Dengue is a global public health problem illness caused by dengue virus and can range from non specific febrile illness to classic dengue fever, which may then progress to dengue haemorrhagic fever and dengue shock syndrome. Average incidence rate is 21 to 50 per million population in Karnataka from 1998 to 2014 [6].

Dengue fever is caused by the dengue virus with one of the four serotypes: DENV-1, DENV-2, DENV-3, and DENV-4 [7]. WHO has estimated that about 50 million patients are infected with dengue fever annually worldwide and 2.5 billion people live in endemic areas of dengue [8]. It is a challenging workup for the clinicians to detect dengue cases in the early stage before the development of severe manifestations. Paired acute and convalescent phase samples are required to diagnose dengue in the first 3 days after symptoms onset. But Serological tests are unreliable during the early course of illness and also for under developed and developing countries, it is difficult to afford PCR testing and rapid diagnostic tests in all the public health centers [9]. It is important to differentiate

dengue fever from other infectious diseases that requires management with specific anti microbial therapy [10]. Specific diagnostic laboratory tests, even when available may not be accurate in early stages of illness. Serological tests for dengue fever and typhus are frequently negative in early stages of illness. Thus, there is still a role for diagnosis based on haematological parameters.

That is why, WHO recently identified among its global research priorities the need for "clinical and laboratory indicators for early dengue" [11].

Multiple clinical and laboratory features can differentiate dengue from other febrile illness. A recent systematic literature review identified multiple clinical and laboratory features that could potentially differentiate dengue from other febrile illness [9].

Dengue diagnosis based on only clinical presentation is challenging and can lead to misdiagnosis also. In study done by Fernandez E et al., shows that less than 50% of the specimens tested in cases of suspected dengue are confirmed to be positive and the rest were negative for dengue test [8]. The present study aimed to identify the

haematological features useful for discriminating dengue from other febrile illnesses and to evaluate the accuracy of these haematological parameters. Logistic regression model is used on haematological parameters to predict laboratory confirmed dengue cases. Using these data, we built and validated logistic regression models to identify haematological parameters to predict laboratory confirmed dengue.

MATERIALS AND METHODS

A retrospective study designed on 340 patients. Of 340 patients, 170 had laboratory evidence for dengue infection while 170 patients had other febrile illness who were admitted with infections like malaria, filariasis, typhoid, influenza, SARS and were negative for dengue test. All these patients complained about fever for 2 to 5 days. All 340 patients were admitted patients in the SDM medical hospital. Non dengue cases consisted of patient with typhoid, influenza, SARS, leptospirosis, malaria, filariasis. Severity of illness consisted of patients with body temperature above 100.4°F for more than two days.

The study period was between January 2017 and July 2017 in Department of Pathology at the SDM medical college and hospital, Dharwad. We have taken ethical committee approval to conduct the study.

Patients presenting with acute onset fever ($\geq 38.0^{\circ}\text{C}$) above the age of 18 years with no other clinically definitive alternative diagnosis were eligible for inclusion in the study.

Exclusion criteria included 1) patients age <18 years
2) lab results for dengue which are indeterminate.

Demographic, clinical and epidemiological information were recorded on a proforma and consent of the patient was taken. A full blood count was performed on anticoagulated whole blood collected at all time points. EDTA blood were analysed to determine the complete blood count using Sysmex XN – 1000. Calibration by internal and external QC controls was also performed on a regular basis.

The following parameters were listed by the haematology analyser-RBC count, Haemoglobin (Hb), Haematocrit (Hct), Platelet count, WBC count, Neutrophil, Monocytes, Lymphocyte and Eosinophils counts, Mean Corpuscular Volume (MCV), Mean Corpuscular Haemoglobin (MCH), and Mean Corpuscular Haemoglobin Concentration (MCHC).

Dengue can be diagnosed in laboratories using different methods. In the present study, Nonstructural protein 1 (NS1) antigen detection was performed from day 0 to day 5, and indirect diagnosis based on the detection of specific anti-dengue immunoglobulin M (IgM) or immunoglobulin G (IgG) antibodies in patients' sera after day 3. If one of the antibodies (either IgG or IgM) is positive, the sample is designated as positive for dengue infection. All these data were retrieved from the patients and the computer records.

STATISTICAL ANALYSIS

Normal distribution of continuous data was determined using the Kolmogorov-Smirnov test. Continuous variables were

categorised following laboratory or usual cut-off values. Cut-off values for the parameters were as follows-WBCs <4000/ μL , Neutrophils <40%, Lymphocytes <10%, Monocytes <2% Eosinophils <1%, RBCs <4 $\times 10^6$ / μL , Hb <11 g/dL and Platelets <1,00,000/ μL were considered lower than cut-off values. Hematocrit >40%, MCV >80 fL, MCH > 25 pg/cell, MCHC > 33 g/dL, were considered higher than cut-off values. Mann-Whitney U test was used to compare the continuous variables [7]. Categorical variables were analysed using the Fisher's exact test or chi-square test. Statistical significance was set at p-value less than 0.05. Variables found to be statistically significant in univariate analysis were entered into multivariate analysis using a logistic regression model to identify independent risk factors for outcomes of interest. All statistical computations were performed using the SPSS version 20.0 software.

RESULTS

The 340 patients who were investigated in this study had a mean age of 26.75 years (SD age 14.36) and in non dengue cases it was 41.66 years (SD age 14.64).

There were more men 101 cases (59.41%) in the dengue group and more men 102 cases (60%) in the Non dengue group. A significant difference relating to gender between the two groups was observed ($p=0.001$).

Out of 170 patients infected with dengue, 54.71% were positive with NS1 antigen, 52.94% were positive with dengue IgM antibody and 44.71% were positive for dengue Ig G antibody.

The following parameters were significantly lower in patients with DF as compared to Non dengue patients (p -value less than 0.05): WBC, Platelets, Neutrophils, Eosinophils. The following parameters were significantly higher in patients with DF as compared to Non dengue cases ($p<0.05$): PCV, Haemoglobin, RBC count, Lymphocytes, Monocytes. The following parameters show no significant difference in patients with DF compared to patients with non dengue cases ($p>0.05$): MCV, MCH, MCH [Table/Fig-1].

Multiple logistic regression analysis of prediction of dengue cases by different parameters was done using independent variables [Table/Fig-2].

The following variables were independently associated with dengue in multivariable analysis: RBC count (OR=0.47, 95% CI 0.22,1.00) WBC count (OR = 3.94, 95% CI 1.91,8.11), Platelet count (OR = 6.94, 95%CI 3.46,13.90) lymphocyte (OR= 0.47, 95% CI 0.22, 1.02), Eosinophils (OR=3.67, 95% CI 1.9,6.89), MCHC (OR = 0.47, 95% CI 0.25, 0.88)

In the present study, multivariate analysis showed two haematological predictors with positive association to confirm dengue: WBC count <4000/cumm and platelet count <1 lakh/cumm. The study also revealed negative association with respect to RBC count <4 million/cumm, Lymphocytes <10%, Eosinophils <2% and MCHC >33 g/dL [Table/Fig-2].

The model had a sensitivity of 77.65% and specificity of 81.76% with positive predictive value of 80.98% and negative predictive value of 78.53%.

Variables	Groups	Mean	SD	Median	p-value	Non dengue vs Dengue
HB g/dL	Non dengue	12.16	2.97	12	0.0001*	p=0.0003*
	Dengue	13.32	2.63	13		
PCV%or HCT	Non-dengue	37.22	8.22	38	0.0001*	p=0.0001*
	Dengue	40.67	8.52	42		
RBC	Non-dengue	4.13	1.16	4	0.0001*	p=0.0001*
	Dengue	4.74	1.08	5		
WBC	Non-dengue	9737.59	4798.05	8720	0.0001*	p=0.0001*
	Dengue	5449.67	3231.87	4540		
Platelets	Non-dengue	276477.65	131975.64	268500	0.0001*	p=0.0001*
	Dengue	114223.53	93663.05	90000		
Neutrophils	Non-dengue	66.16	15.84	66	0.0001*	p=0.0001*
	Dengue	56.15	20.47	58		
Lymphocytes	Non-dengue	23.84	13.04	24	0.0001*	p=0.0001*
	Dengue	33.66	19.92	32		
Monocytes	Non-dengue	7.81	4.24	7	0.1260	p=0.0479*
	Dengue	8.45	4.27	8		
Eosinophils	Non-dengue	2.12	3.51	1	0.0001*	p=0.0001*
	Dengue	1.14	2.16	0		
MCV	Non-dengue	85.81	9.59	86	0.5570	p=0.6125
	Dengue	85.60	6.98	86		
MCH	Non-dengue	27.95	3.74	29	0.3060	p=0.0952
	Dengue	27.68	3.10	28		
MCHC	Non-dengue	32.34	2.01	32	0.0030*	p=0.5706
	Dengue	39.39	46.00	32		

[Table/Fig-1]: Pair wise comparisons by Mann-Whitney U test

Independent variables	Odds Ratio	p-value	95% CI for OR	
HB <11g/dl	0.53	0.1680	0.21	1.31
PCV% or HCT>40%	0.97	0.9380	0.50	1.89
RBC<4x10 ⁶ /uL	0.47	0.0500*	0.22	1.00
WBC<4000/ μ L	3.94	0.0001*	1.91	8.11
Platelets<100000/ μ L	6.94	0.0001*	3.46	13.90
Neutrophils<2800/ μ L	1.56	0.3460	0.62	3.95
Lymphocytes<800/ μ L	0.47	0.0500*	0.22	1.02
Monocytes<80/ μ L	1.18	0.7980	0.34	4.15
Eosinophils<40/ μ L	3.67	0.0001*	1.97	6.83
MCV>80fL	1.01	0.9790	0.38	2.72
MCH>25pg/cell	1.06	0.9110	0.36	3.10
MCHC>33g/dL	0.47	0.0190*	0.25	0.88

[Table/Fig-2]: Multiple logistic regression analysis of prediction of dengue cases by different parameters.
*p<0.05

DISCUSSION

The present study confirmed that socio-demographic characteristics differ between patients with DF and those with non dengue fever patients.

Firstly, patients with DF tended to be younger than non dengue patients, however, no significant difference between the two groups and age were observed (mean age DF = 26.75, ND = 41.66 years old). This was in concordance to another study conducted by Kotepui M et al., which, indicated that patients with DF were younger [12]. Another study done by Gregory CJ et al., showed that laboratory positive dengue cases were older than patients who were laboratory negative for dengue [9].

Secondly, males were infected more in dengue cases in the present study. This was in concordance with the study done by Fernandez E et al., and Daumas P et al., in which males accounted for 59% and 65.1% in cases respectively [8,13].

However, in study done by Hammond SN et al., females were more affected than males [14]. In the present study, haematological parameters were more severe and

abnormalities were more frequent in patient with dengue than those with non dengue.

The following parameters were significantly lower in patients with DF as compared to non dengue patients (p-value less than 0.05): WBC, Platelets, Neutrophils, Eosinophils. Transient marrow suppression is the cause of Neutropenia in DF [15]. Platelet survival time is shortened due to multiple mechanisms leading to platelet destruction [15]. In initial stages of dengue illness there is hypocellularity of bone marrow and attenuation of megakaryocytes maturation [16].

In study done by Kalyanarooj S et al., Platelet count, Total WBC, Neutrophils and Monocytes were significantly lower in DF than OFI [17].

The following parameters were significantly higher in patients with DF as compared to non dengue cases (p<0.05): PCV, Haemoglobin, RBC count, Lymphocytes, Monocytes. Higher values of PCV vascular leakages are one of the causes for increase in PCV leading to haemo concentration [18]. One study showed that lymphocyte count was within normal limits during the course of dengue illness [16]

Rising values of hematocrit and rapid decrease of platelet count indicates plasma leakage in dengue patient [19].

In study done by Kotepui M et al., RBC, Hb, PCV, MCV, MCH, MCHC were significantly higher in patients with dengue. Parameters with significantly lower values were WBC, Neutrophils, Monocytes, Eosinophils. In another study done by Dumas P et al., Platelet and Leucocytes counts were significantly lower in dengue than in non dengue groups [13].

Our logistic regression model found that five haematological parameters helped differentiate dengue from other febrile illness:

- 1) WBC count;
- 2) Platelet count;
- 3) RBC count;
- 4) Lymphocytes;
- 5) Eosinophils.

Of these, the association was positive for the WBC count and Platelet count (likely to present in dengue cases) and association was negative for the RBC count, Lymphocytes, Eosinophils, MCHC consistent with being less likely to be present in dengue cases.

Platelet count has a higher odds ratio than other predictors followed by WBC count. Therefore, low platelet count and low WBC count are strong positive predictors in dengue cases. In study done by Gregory CJ et al., low platelet count had a odds ratio of 2.1 (95% CI 1.3, 3.4) [9]

Hammond SN et al., showed that platelet count less than 150,000/mm³, 100,000/mm³ and 50,000/mm³ having positive laboratory diagnosis for dengue. Platelet was significantly lower in DF than OFI and Hematocrit were significantly higher in DF than in OFI [14].

The platelet cut-off of <140,000/mm³ is shown to be predictors in distinguishing dengue from other febrile illnesses [20].

One study showed that Leucopenia, Neutropenia, Monocytopenia and increased AST levels can be used as predictors of dengue in early stage before plasma leakage

developed thus help in close monitoring of patient [17]. This study also showed that Hematocrit and Thrombocytopenia are critical features that distinguish DF from DHF [17].

In a study done by Wilder Smith A et al., the multivariate analysis showed that predictors of dengue from other febrile illnesses were low platelet count and low WBC count and elevated AST level and these three parameters had high odds ratio [21].

In a study done by Low LG et al., platelet count was significantly lower in DF than OFI. And lymphocyte counts were significantly higher. But Neutrophil, PCV, WBC were not significant [22].

However, Deparis X et al., mentions in his study that these laboratory measures are not specific for dengue in early stage of disease as the study showed low odds ratio for the parameters [23].

Chadwick D et al., showed that WBC count lower than 5000/cumm was the only laboratory parameter which was predictive of dengue fever [24].

In a study done by La Russa VF et al., showed that dengue infected bone marrow stromal cells and dengue specific T cells produce few cytokines which cause bone marrow suppression [25].

LIMITATION

The limitations of the study are, since it is a retrospective study, we could not assess the variation in the clinical and laboratory features that took place during the course of illness. Secondly the data is from only one region that is north Karnataka region. So generalisation of results is difficult and samples from other parts of the state need to be studied. Thirdly, serologically negative dengue cases were not considered in dengue case group as we considered sero positivity as diagnostic criteria for dengue case.

This study suggests that simple laboratory investigation like CBC can be used to identify early dengue infection from other febrile illness in adults.

The study implicates that an algorithm is needed to identify the patients with dengue early in illness. So that unnecessary hospitalisation is avoided.

The study highlights that many more prospective studies are required so as to set up a clinical and laboratory algorithm that can validate and generalise between dengue fever and other febrile illnesses.

The study broadens the horizon in the literature of dengue illness that leucopenia especially affecting Neutrophils and Monocyte lineage is helpful to differentiate dengue from other febrile illnesses.

CONCLUSION

Dengue will continue to increase and spread day by day until a safe and effective vaccine is available and a viable and unending mosquito control practice takes place.

This study helps the clinicians to make an empirical diagnosis when diagnostic tools and efficient trained personnel are not

available. And also helps to start treatment in emergency cases of these diseases when the reports are awaited in endemic areas.

The present study also finds a way to avoid a large number of other investigations to diagnose dengue from other febrile illnesses. However, the rising availability of rapid tests for dengue and other illness may diminish the interest of platelet count and WBC count to differentiate febrile illnesses.

REFERENCES

- [1] Epelboin L, Boullé C, Ouar-Epelboin S, Hanf M, Dussart P, Djossou F, et al. Discriminating malaria from dengue fever in endemic areas: clinical and biological criteria, prognostic score and utility of the C-reactive protein: a retrospective matched-pair study in French Guiana. *PLoS Negl Trop Dis*. 2013;7(9):e2420.
- [2] Simmons C P, J. Farrar J, VinhChau N, and Wills B. Current concepts: dengue. *The New England Journal of Medicine*. 2012;366(15):1423-32.
- [3] Anuradha S, Singh NP, Rizvi SN, Agarwal SK, Gur R, Mathur MD. The 1996 outbreak of dengue hemorrhagic fever in Delhi, India. *Southeast Asian J Trop Med Public Health*. 1998;29(3):503-06.
- [4] Ralapanawa DMPUK, Jayawickreme KP, Ekanayake EMM, Jayalath T, Herath D. Fatal massive pulmonary hemorrhage in dengue infection. *Epidemiology*. 2016;6(251):2161-65.
- [5] Ralapanawa DM, Kularatne SA, Jayalath WA. Guillain-Barre syndrome following dengue fever and literature review. *BMC Res Notes*. 2015;8(1):729.
- [6] Mutheneeni S R, Morse A P, Caminade C and Upadhyayula S M. Dengue burden in India: recent trends and importance of climatic parameters. *Emerging Microbes & Infections*. 2017;6(8):57-58.
- [7] Kotepui M, PhunPhuech B, Phiwklam N, Uthaisar K.. Differentiating between dengue fever and malaria using haematological parameters in endemic areas of Thailand. *Infect Dis Poverty*. 2017;6(1):27.
- [8] Fernández E, Smieja M, Walter SD, Loeb M. A predictive model to differentiate dengue from other febrile illness. *BMC Infect Dis*. 2016; 16: 694.
- [9] Gregory C.J, Santiago LM, Argüello DF, Hunsperger E, Tomashek KM. Clinical and laboratory features that differentiate dengue from other febrile illnesses in an endemic area—Puerto Rico. *Am J Trop Med Hyg*. 2010;82(5):922-29.
- [10] Bruce MG, Sanders EJ, Leake JA, Zaidel O, Bragg SL, Aye T, et al. Leptospirosis among patients presenting with dengue-like illness in Puerto Rico. *Acta Trop*. 2005;96(1):36-46.
- [11] Dengue haemorrhagic fever: diagnosis, treatment, prevention and control. 2nd edition. Geneva : World Health Organization. 1997. <http://www.who.int/csr/resources/publications/dengue/Denguepublication/en/>
- [12] Kotepui M, Phunphuech B, Phiwklam N, Chupeerach C, Duangmano S. Effect of malarial infection on haematological parameters in population near Thailand-Myanmar border. *Malar J*. 2014; 13: 218.
- [13] Daumas P, Passos Sonia RL , Oliveira Raquel VC , Nogueira Rita MR , Georg I, Marzochi Keyla BF, et al. Clinical and laboratory features that discriminate dengue from other febrile illnesses: a diagnostic accuracy study in Rio de Janeiro, Brazil. *BMC Infectious Diseases*. 2013;13:77
- [14] Hammond SN, Balmaseda A, Pérez L, Tellez Y, Saborío SI, Mercado JC, et al. Differences in dengue severity in infants, children, and adults in a 3-year hospital-based study in Nicaragua. *Am J Trop Med Hyg*. 2005;73(6):1063-70.
- [15] Potts JA, Rothman AL. Clinical and laboratory features that distinguish dengue from other febrile illnesses in endemic population. *Trop Med Int Health*. 2008;13(11):1328-40.
- [16] Lam PK, Ngoc TV, Thu Thuy TT, Hong Van NT, Nhu Thuy TT, Hoai Tam DT, et al. The value of daily platelet counts for predicting dengue shock syndrome: results from a prospective observational study of 2301 Vietnamese children with dengue. *PLoS Negl Trop Dis*. 2017 Apr 27;11(4):e0005498.
- [17] Kalayanarooj S, Vaughn DW, Nimmanitya S, Green S, Suntayakorn S, Kunentrasai N, et al. Early clinical and laboratory indicators of acute dengue illness. *J Infect Dis*. 1997;176(2):313-21.
- [18] Cardier JE, Mariño E, Romano E, Taylor P, Liprandi F, Bosch N, et al. Proinflammatory factors present in sera from patients with acute dengue infection induce activation and apoptosis of human microvascular endothelial cells: possible role of TNF-alpha in endothelial cell damage in dengue. *Cytokine*. 2005 Jun 21;30(6):359-65.
- [19] Ralapanawa U, Alawattegama ATM, Gunrathne M, Tennakoon S, Kularatne SAM, Jayalath T. Value of peripheral blood count for dengue severity prediction. *BMC Res Notes*. 2018;11(1):400.
- [20] Watt G, Jongsakul K, Chouriyagune C, Paris R. Differentiating dengue virus infection from scrub typhus in Thai adults with fever. *Am J Trop Med Hyg*. 2003 May;68(5):536-38.
- [21] Wilder-Smith A, Earnest A, Paton NI. Use of simple laboratory features to distinguish the early stage of severe acute respiratory syndrome from dengue fever. *Clin Infect Dis*. 2004;39:1818-23.
- [22] Low JG , Ooi EE, Tolfvenstam T, Leo YS, Hibberd ML, Ng LC, et al. Early Dengue infection and outcome study (EDEN)-study design and preliminary findings. *Ann Acad Med Singapore*. 2006 ;35(11):783-89.
- [23] Deparis X , Murgue B, Roche C, Cassar O, Chungue E. Changing clinical and biological manifestations of dengue during the dengue-2 epidemic in French Polynesia in 1996/97 -description and analysis in a prospective study. *Trop Med Int Health*. 1998;3(11):859-65.
- [24] Chadwick D , Arch B, Wilder-Smith A, Paton N. Distinguishing dengue fever from other infections on the basis of simple clinical and laboratory features: application of logistic regression analysis. *J Clin Virol*. 2006;35(2):147-53.
- [25] La Russa VF, Innis BL. Mechanisms of dengue virus-induced bone marrow suppression. *Baillieres Clin Haematol*. 1995;8(1):249-70.

AUTHOR(S):

1. Dr. Priyanka P
2. Dr. Dinesh US

PARTICULARS OF CONTRIBUTORS:

1. Assistant Professor, Department of Pathology, SDM College of Medical Sciences and Hospital, Dharwad, Karnataka, India.
2. Professor, Department of Pathology, SDM College of Medical Sciences and Hospital, Dharwad, Karnataka, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Priyanka P,
Assistant Professor, Department of Pathology, SDM
College of Medical Sciences and Hospital, Dharwad,
Karnataka, India.
E-mail: drpriyankamahesh@gmail.com

FINANCIAL OR OTHER COMPETING INTERESTS:

None.

Date of Publishing: Oct 01, 2018